

VINBERG'S θ -GROUPS AND RIGID CONNECTIONS

TSAO-HSIEN CHEN

ABSTRACT. Let G be a simple complex group of adjoint type. In his unpublished work, Z. Yun associated to each θ -group $(G_0, \mathfrak{g}_1)$ and a vector $X \in \mathfrak{g}_1$ a flat G -connection ∇^X on $\mathbb{P}^1 - \{0, \infty\}$, generalizing the construction of Frenkel and Gross in [FG]. In this paper we study the local monodromy of those flat G -connections and compute the de Rham cohomology of ∇^X with values in the adjoint representations of G . In particular, we show that in many cases the connection ∇^X is cohomologically rigid.

CONTENTS

1. Introduction	1
2. Gradings on simple Lie algebras	3
3. Kac-Moy-Prasad gradings on loop algebras	6
4. Yun's θ -connections	9
5. Main result	11
6. Proofs	12
7. Examples	15
8. Relation with [Yun] and ramified geometric Langlands	18
References	18

1. INTRODUCTION

1.1. **The goal.** Let G be a simple complex algebraic group of adjoint type. Motivated by Langlands correspondence, Frenkel and Gross [FG] constructed a flat G -connection ∇ on the trivial G -bundle on $\mathbb{P}^1 - \{0, \infty\}$ with following remarkable properties:

2010 *Mathematics Subject Classification.* 17B67, 22E50, 22E57.

- (1) ∇ has regular singularity at 0, and the residue is a regular nilpotent element in the Lie algebra $\mathfrak{g}$ of G .
- (2) ∇ has irregular singularity at ∞ with slope $1/h$, where h is the Coxeter number of G (see [FG, §5] or [CK, §2.2]) for the definition of slope).
- (3) ∇ is *cohomologically rigid*, i.e., we have $H^*(\mathbb{P}^1, j_{!*}\nabla^{\text{Ad}}) = 0$, here ∇^{Ad} is the D -module defined by the connection ∇ with values in the adjoint representation of G and $j_{!*}\nabla^{\text{Ad}}$ is the intermediate extension of the D -module ∇^{Ad} to $\mathbb{P}^1$.

The construction used the θ -group $(G_0, \mathfrak{g}_1)$ studied by Vinberg and his school, which comes from a $\mathbb{Z}/h\mathbb{Z}$ -grading on $\mathfrak{g}$.

In his unpublished work [Yun1], Yun generalized the Frenkel-Gross's construction to *all* θ -groups. More precisely, starting with a θ -group $(G_0, \mathfrak{g}_0)$, he constructed a family of flat G -connections ∇^X on $\mathbb{P}^1 - \{0, \infty\}$ parametrizing by vectors $X \in \mathfrak{g}_0$. We called ∇^X the θ -connection associated to $X \in \mathfrak{g}_1$.

The goal of this paper is to study properties of the θ -connections ∇^X . In more detail, recall that each θ -group $(G_0, \mathfrak{g}_1)$ corresponds to a torsion automorphism of $\mathfrak{g} = \text{Lie}G$, which we also denote by $\theta \in \text{Aut}(\mathfrak{g})$. Let σ be the image of θ in $\text{Out}(\mathfrak{g})$, the group of outer automorphism of $\mathfrak{g}$. We establish the following properties of ∇^X , parallel to the properties (1), (2) and (3) above:

- (1) ∇^X has regular singularity at 0, and the residue $\text{Res}(\nabla^X)$ is a nilpotent element in the Lie algebra $\mathfrak{g}^\sigma$. Moreover, for generic $X \in \mathfrak{g}_1$ the residue $\text{Res}(\nabla^X)$ lies in a single nilpotent orbits in $\mathfrak{g}^\sigma$.
- (2) ∇^X has irregular singularity at ∞ with slope e/m for any semi-simple $X \in \mathfrak{g}_1$. Here m (resp. e) is the order of $\theta \in \text{Aut}(\mathfrak{g})$ (resp. $\sigma \in \text{Out}(\mathfrak{g})$).
- (3) Assume θ is *stable* with normalized *Kac coordinates* $s_0 = 1$ (see §2.2 for the definition of stable automorphism and normalized Kac coordinates). Then for any stable element $X \in \mathfrak{g}_1$ the connection ∇^X is *cohomologically rigid*.

1.2. Let me explain briefly how these properties are obtained. Properties (1) and (2) follow basically from the construction. To prove property (3), we need to show that certain cohomology groups vanish (see Definition 4.1). Here we follow an argument in [FG]. First, it follows from a general result in [FG, §7] that the relevant cohomology groups are isomorphic to kernels of certain $\mathbb{C}$ -linear maps on the loop algebra $\tilde{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}]$ (or its twisted version ${}^\sigma\tilde{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}]^\sigma$). So we reduce to show that kernels of those maps vanish. In the case of [FG], the authors observe that if we consider the *principal* grading on the loop algebra $\tilde{\mathfrak{g}}$ (see Example 3.1), then in terms of a homogeneous basis of $\tilde{\mathfrak{g}}$ (with respect to the principal grading), the relevant maps become more tractable and the desired cohomology vanishing follows from Kac's Theorem on *principal* Heisenberg subalgebras of affine Kac-Moody algebras. Now our observation is that for general θ -connections ∇^X , if we consider the *Kac-Moy-Prasad* grading on $\tilde{\mathfrak{g}}$ corresponding to the torsion automorphism θ (see §3), then the relevant maps again become more tractable and the desired cohomology vanishing

follows from the results in [Kac, RLYG, RY] about gradings on Lie algebras and loop algebras, and a generalization of Kac's Theorem to *general* Heisenberg subalgebras of affine Kac-Moody algebras (see Proposition 3.5 and Remark 4.2).

1.3. Relation with [Yun]. In [Yun], starting with a stable torsion automorphism $\theta \in \text{Aut}(\check{\mathfrak{g}})$ for the Langlands dual of $\mathfrak{g}$ and a stable functional $\phi \in \check{\mathfrak{g}}_1^{*,s}$, the author construct a remarkable ℓ -adic G -local system $\text{KL}_G(\phi)$ on $\mathbb{P}^1 - \{0, \infty\}$, which generalized his early work in [HNY] with Heinloth and Ngô. This ℓ -adic local system is tamely ramified at 0 and ramified at ∞ . He furthermore describe the monodromy of $\text{KL}_G(\phi)$ at 0 and conditionally deduce the cohomologically rigid of $\text{KL}_G(\phi)$ (cf. [Yun, Theorem 4.7 and Proposition 5.2]). The construction can carry out over the complex number with ℓ -adic sheaves replaced by D -modules. Thus, starting with a stable automorphism θ of $\check{\mathfrak{g}}$ and a stable function $\phi \in \check{\mathfrak{g}}_1^{*,s}$, we get a flat G -connection $\text{KL}_G(\phi)_{dR}$ on $\mathbb{P}^1 - \{0, \infty\}$. The result of this paper gives strong evidence that the θ -connection ∇^X is isomorphic to $\text{KL}_G(\phi)_{dR}$ (see Conjecture 8.1 in §8). In the case when θ has order h , this is proved in [Zhu].

1.4. The paper is organized as follows. In §2 we give a review of Vinberg's theory of θ -groups. In §3 we recall the Kac-Moy-Prasad grading for loop algebras and Kac's theories on automorphism of loop algebras and Heisenberg subalgebras of affine Kac-Moody algebras. In §4 we recall Yun's construction of θ -connections ∇^X . We compute the residue of ∇^X at 0 and the slope and irregularity at ∞ . In §5 we prove the main results of this paper: In Theorem 5.1, assume the θ -group $(G_0, \mathfrak{g}_1)$ is regular, i.e., $\mathfrak{g}_1$ contains regular semi-simple elements, we compute the de Rham cohomology of the intermediate extension to $\mathbb{P}^1$ of the D -module on $\mathbb{P}^1 - \{0, \infty\}$ defined by the θ -connection ∇^X associated to a regular semi-simple $X \in \mathfrak{g}_1$ with values in the adjoint representation of G . In Theorem 5.2, assume the θ -group is stable and with normalized Kac coordinates $s_0 = 1$, we establish the cohomological rigidity of ∇^X . In §7 we give several examples of θ -connections. Finally, in §8 we discuss the relation with [Yun] and ramified geometric Langlands.

Acknowledgement. The author is grateful to Z. Yun for allowing him to use his unpublished result and for many helpful discussions. An ongoing project with M. Kamgarpour and X. Zhu is the main motivation to write this paper. The author thanks them cordially for the collaboration. The main part of this work was accomplished when the author was visiting the Max Planck Institute for Mathematics in Bonn. He thank the institution for the wonderful working atmosphere.

2. GRADINGS ON SIMPLE LIE ALGEBRAS

2.1. Notation. Let $\mathfrak{g}$ be the Lie algebra of a simple complex algebraic group G of adjoint type. Let ℓ be the rank of $\mathfrak{g}$. Let B be a Borel subgroup of G and let $T \subset B$ be maximal torus. We denote by $\mathfrak{g} = \text{Lie}G$, $\mathfrak{t} = \text{Lie}T$ and $\mathfrak{b} = \text{Lie}B$. We let

$\text{Ad} : G \rightarrow \text{Aut}(\mathfrak{g})$ (resp. $\text{ad} : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g})$) denote the adjoint representation of G (resp. $\mathfrak{g}$). For any element $x \in \mathfrak{g}$ we denote by $\mathfrak{g}^x$ the kernel of $\text{ad}(x)$.

Let $\mathbb{X}$ (resp. $\check{\mathbb{X}}$) be the weight lattices (resp. coweight lattices) of T , and R (resp. $\check{R}$) be the set of roots (resp. co-roots) of T in G . We fixed a pinning $(\mathbb{X}, R, \check{\mathbb{X}}, \check{R}, \{E_i\})$, where $E_i \in \mathfrak{g}$ is a root vector for the simple roots $\alpha_i \in \Delta$.

Let $\text{Aut}(R)$ be the subgroup of $\text{Aut}(\mathbb{X})$ preserving R and let $\text{Aut}(R, \Delta)$ be the subgroup of $\text{Aut}(R)$ preserving Δ . The choice of pinning induced an isomorphism $\text{Aut}(\mathfrak{g}) = G \rtimes \text{Aut}(R, \Delta)$. Let $\exp : V := \check{\mathbb{X}} \otimes \mathbb{R} \rightarrow T$ be the exponential map given by $\alpha(\exp(x)) = e^{2\pi i \alpha(x)}$, for all $\alpha \in \mathbb{X}$.

For any $m \in \mathbb{Z}^\times$, we let $\xi_m = e^{\frac{2\pi i}{m}}$. For any $\mathbb{C}$ -vector space V , we denote by $V^* = \text{Hom}(V, \mathbb{C})$ the dual of V .

2.2. Affine simple roots. In this subsection we collect some basic definitions and properties of twisted affine Kac-Moody algebra and affine simple roots. For details, see [Kac, §8].

Let $\sigma \in \text{Aut}(R, \Delta)$ and let e be the order of σ . We have $e = 1$ or $e = 2$ (type A, D, E_6) or $e = 3$ (type D_4). Consider the affine Kac-Moody algebra $\hat{\mathfrak{g}} = \tilde{\mathfrak{g}} \oplus \mathbb{C}K \oplus \mathbb{C}d$. Here $\tilde{\mathfrak{g}} \oplus \mathbb{C}K$ is the universal central extension of the loop algebra $\tilde{\mathfrak{g}} := \mathfrak{g}[t, t^{-1}]$ and $[d, t^i \otimes x + K] = it^i \otimes x$. The automorphism σ extends to an automorphism of $\hat{\mathfrak{g}}$ by

$$\sigma(t^i \otimes x) = \xi_e^{-i} t^i \otimes \sigma(x), \quad \sigma(K) = K, \quad \sigma(d) = d.$$

The fixed point subalgebra

$${}^\sigma \hat{\mathfrak{g}} := {}^\sigma \tilde{\mathfrak{g}} \oplus \mathbb{C}K \oplus \mathbb{C}d,$$

where ${}^\sigma \tilde{\mathfrak{g}} := \mathfrak{g}[t, t^{-1}]^\sigma$, is the twisted affine Kac-Moody algebra associated to $(\mathfrak{g}, \sigma)$.

Set ${}^\sigma \hat{\mathfrak{t}} := \mathfrak{t}^\sigma \oplus \mathbb{C}K \oplus \mathbb{C}d$ (here $\mathfrak{t}^\sigma$ is the σ -fixed vectors in $\mathfrak{t}$) and define $\delta \in ({}^\sigma \hat{\mathfrak{t}})^*$ by $\delta|_{\mathfrak{t}^\sigma \oplus \mathbb{C}K} = 0$, $\delta(d) = 1$. Then there is an affine root spaces decomposition of ${}^\sigma \hat{\mathfrak{g}}$ with respect to ${}^\sigma \hat{\mathfrak{t}}$

$$(1) \quad {}^\sigma \hat{\mathfrak{g}} = {}^\sigma \hat{\mathfrak{t}} \oplus \bigoplus_{\alpha \in \Phi_{\text{aff}}} {}^\sigma \hat{\mathfrak{g}}_\alpha$$

here $\Phi_{\text{aff}}^\sigma \subset (\mathfrak{t}^\sigma)^* \oplus \mathbb{C}\delta$ is the set of affine roots. We identify affine roots Φ_{aff}^σ with affine functions on $\mathfrak{t}^\sigma$ by sending δ to the constant function 1.

We now recall the construction of affine simple roots $\Delta_{\text{aff}}^\sigma \subset \Phi_{\text{aff}}^\sigma$. Let R/σ (resp. Δ/σ) be the set of orbits in R (resp. Δ) under σ . For any orbit $a \in R/\sigma$, let β_a denote the restriction to $\mathfrak{t}^\sigma$ of any $\alpha \in a$. Then the collection $R^\sigma := \{\beta_a | a \in R/\sigma\}$ is a root system (possibly non-reduced) with basis $\Delta^\sigma = \{\beta_a | a \in \Delta/\sigma\}$. Let us choose a numbering

$$\Delta^\sigma = \{\beta_1, \dots, \beta_{\ell_\sigma}\},$$

where ℓ_σ is the number of σ -orbits on the set of simple roots Δ . We define a certain positive root η as follows. If $\sigma = id$, then η is the highest root. If $\sigma \neq id$, unless (R, σ) is of type ${}^2A_{2n}$, η is the highest short root of R^σ ; when (R, σ) is of type ${}^2A_{2n}$, η is twice the highest short root of R^σ . There are unique positive integers $\{b_0 = 1, b_1, \dots, b_{\ell_\sigma}\}$ such that

$$(2) \quad \eta = \sum_{i=1}^{\ell_\sigma} b_i \beta_i.$$

The sum $h_\sigma := e \sum_{i=0}^{\ell_\sigma} b_i$ is the twisted Coxeter number of (R, σ) . We define

$$\beta_0 = 1/e - \eta.$$

Then $\Delta_{\text{aff}}^\sigma := \{\beta_0, \dots, \beta_{\ell_\sigma}\}$ is the set of affine simple roots associated to $(\mathfrak{g}, \sigma)$. The set $C^\sigma = \{x \in V^\sigma \mid \beta_i(x) > 0, i = 0, \dots, \ell_\sigma\}$ is called the fundamental alcove. We denote by $\bar{C}^\sigma$ the closure of C^σ in V^σ .

2.3. Normalized Kac coordinates. Let $\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}/m\mathbb{Z}} \mathfrak{g}_i$ be a grading of $\mathfrak{g}$. The grading on $\mathfrak{g}$ corresponds to a torsion automorphism $\theta = \theta' \rtimes \sigma \in \text{Aut}(\mathfrak{g})$ such that $\theta(v) = \xi_m^i v$. The automorphism θ is G -conjugate to one of the form $t \rtimes \sigma$ with $t \in T^\sigma$. Thus without loss of generality, we can assume $\theta = t \rtimes \sigma$, $t \in T^\sigma$.

According to [Kac, Proposition 8.1] (see also [OV, §3]), there exists $x \in \bar{C}^\sigma$ such that $\theta = \exp(x) \rtimes \sigma$. Since θ has order m , we have

$$(3) \quad \beta_i(x) = \frac{s_i}{m}.$$

The integers $(s_i)_{i=0, \dots, \ell_\sigma}$ are the normalized *Kac coordinates* of θ (cf. [RLYG, §2.2]). These coordinates satisfy

$$(4) \quad e \sum_{i=0}^{\ell_\sigma} b_i s_i = m,$$

here e is the order of σ and the b_i are integers mentioned earlier in §2.2.

Let $\check{\lambda} = mx \in \check{X}^\sigma$. The action of $\mathbb{G}_m$ on $\mathfrak{g}$ via $\check{\lambda}$ give a grading $\mathfrak{g} = \bigoplus_{k \in \mathbb{Z}} \mathfrak{g}(k)$ and each $\mathfrak{g}_i$ decomposes as

$$(5) \quad \mathfrak{g}_i = \bigoplus_{k \in \mathbb{Z}} \mathfrak{g}_i(k).$$

Lemma 2.1. *In above decomposition, we have 1) $\mathfrak{g}_i(k) = 0$ unless $k \equiv i \pmod{\frac{m}{e}}$ and 2) $-m + es_0 \leq k \leq m - es_0$.*

Proof. Let $0 \neq v \in \mathfrak{g}_i(k)$. Then we have $\theta(v) = \xi_m^i v$. On the other hand, since $\theta = \check{\lambda}(\xi_m) \rtimes \sigma$, we have $\theta(v) = \sigma(\xi_m^k v) = \xi_e^l \xi_m^k v = (\xi_m)^{\frac{ml}{e} + k} v$ for some $l \in \mathbb{Z}$. This implies $k \equiv i \pmod{\frac{m}{e}}$. For part 2), it is enough to show that $\beta(\check{\lambda}) \leq m - es_0$, where

β is the highest root of $\mathfrak{g}$ (here we regard β as an element in $(\mathfrak{t}^\sigma)^*$). For this, observe that we have $e\eta - \beta \in \sum_{i=1}^{\ell_\sigma} \mathbb{Z}_{\geq 0} \beta_i^1$, here η is the root introduced in (2). Hence

$$\beta(\check{\lambda}) \leq e\eta(\check{\lambda}) = e \sum_{i=1}^{\ell_\sigma} b_i \beta_i(\check{\lambda}) \stackrel{(3)}{=} e \sum_{i=1}^{\ell_\sigma} b_i s_i \stackrel{(4)}{=} m - es_0.$$

We are done. □

2.4. The θ -group. Let G_0 be the reductive subgroup of G with Lie algebra $\mathfrak{g}_0$. There are natural actions of $G_0 \rtimes \sigma$ on $\mathfrak{g}_i$. The pair $(G_0, \mathfrak{g}_1)$ is called θ -group in the terminology of the Vinberg school.

A grading $\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}/m\mathbb{Z}} \mathfrak{g}_i$ (resp. a torsion automorphism $\theta \in \text{Aut}(\mathfrak{g})$, resp. a θ -group $(G_0, \mathfrak{g}_1)$) is called *regular*, if $\mathfrak{g}_1$ contains a regular semi-simple element; *stable* if $\mathfrak{g}_1$ contains a stable element (recall that an element v is called stable if G_0 -orbit of v is closed and the stabilizer in G_0 is finite). According to [RLYG, §5.3], a vector $v \in \mathfrak{g}_1$ is stable if only if v is a regular semi-simple elements of $\mathfrak{g}$ and the action of θ on the Cartan sub-algebra centralizing v is *elliptic*, i.e. $Z_{\mathfrak{g}_0}(v) = 0$. We denote by $\mathfrak{g}_1^r$ (resp. $\mathfrak{g}_1^s$) the open set of regular semi-simple (resp. stable) elements.

For future reference, we include a lemma about structure of $\mathfrak{g}_0$:

Lemma 2.2. 1) $\mathfrak{g}_0$ is the reductive subalgebra with Cartan subalgebra $\mathfrak{t}^\sigma$ and the system of simple roots $\Delta_0^\sigma = \{\bar{\beta}_i | \beta_i \in \Delta_{\text{aff}}^\sigma, s_i = 0\}$. Here $\bar{\beta}_i$ denotes the linear part of β_i . 2) If $s_0 \neq 0$, we have $\mathfrak{g}_0 = \mathfrak{g}_0 \cap \mathfrak{g}(0) = \mathfrak{g}(0)^\sigma$.

Proof. Part 1) is proved in [Kac, Proposition 8.6] (see also [OV, §3.11]). Part 2) follows from part 1) together with the fact $\mathfrak{g}(0)^\sigma$ is the Levi subalgebra of $\mathfrak{g}^\sigma$ with system of simple roots $\{\bar{\beta}_i | \beta_i \in \Delta_{\text{aff}}^\sigma, i \neq 0, s_i = 0\}$. □

3. KAC-MOY-PRASAD GRADINGS ON LOOP ALGEBRAS

3.1. Let ${}^\sigma\tilde{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}]^\sigma$ be the twisted loop algebra. The decomposition in (1) induces a roots space decomposition of the twisted loop algebra ${}^\sigma\tilde{\mathfrak{g}} = \bigoplus_{\alpha \in \Phi_{\text{aff}}^\sigma \cup \{0\}} {}^\sigma\tilde{\mathfrak{g}}_\alpha$ (here we set ${}^\sigma\tilde{\mathfrak{g}}_0 = \mathfrak{t}^\sigma$). For any $x \in V^\sigma$, Kac, Moy and Prasad have introduced a $\mathbb{Z}$ -grading

$$(6) \quad {}^\sigma\tilde{\mathfrak{g}} = \bigoplus_{i \in \mathbb{Z}} {}^\sigma\tilde{\mathfrak{g}}_{x,i}.$$

Explicitly, we have

$${}^\sigma\tilde{\mathfrak{g}}_{x,i} = \bigoplus_{\alpha \in \Phi_{\text{aff}}^\sigma \cup \{0\}, \alpha(x) = \frac{i}{m}} {}^\sigma\tilde{\mathfrak{g}}_\alpha.$$

¹This is obvious when $\sigma = id$, for $\sigma \neq id$ one can use the table 1 in [RLYG] to check it.

We called the $\mathbb{Z}$ -grading in (6) the *Kac-Moy-Prasad* grading associated to x .

Example 3.1 (Principal grading). Consider the case $\theta = \exp(x)$ where $x = \frac{\check{\rho}}{h}$, $\check{\rho}(\alpha_i) = 1$ for $i = 1, \dots, \ell$. In this case the corresponding Kac-Moy-Prasad grading can be described as follows. Let $E_i, H_i, F_i, i = 1, \dots, \ell$ be a Chevalley basis of $\mathfrak{g}$ such that $E_i \in \mathfrak{n}, H_i \in \mathfrak{t}, F_i \in \bar{\mathfrak{n}}$. Then $\tilde{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}]$ has Kac-Moody generators $e_i = E_i \otimes 1, h_i = H_i \otimes 1, f_i = F_i \otimes 1, i = 1, \dots, \ell; e_0 = F_{-\beta} \otimes t, f_0 = E_{\beta} \otimes t^{-1}$, here β is the highest root and E_{β} (resp. $F_{-\beta}$) is a generator of the root space $\mathfrak{g}_{\beta}$ (resp. $\mathfrak{g}_{-\beta}$). Now the grading on $\tilde{\mathfrak{g}}$ is given by $\deg(e_i) = -\deg(f_i) = 1, \deg(h_i) = 0$. We call this grading the *principal* grading (see [Kac, §14]).

Remark 3.2. In [Kac, §1.3], Kac called the $\mathbb{Z}$ -grading in (6) the gradation of type $(s_0, \dots, s_{\ell_{\sigma}})$.

3.2. We preserve the setup in §2.3. Let $\theta = \exp(x) \rtimes \sigma \in \text{Aut}(\mathfrak{g}), x \in \bar{C}^{\sigma}$ and let $\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}/m\mathbb{Z}} \mathfrak{g}_i$ be the grading on $\mathfrak{g}$ defined by the automorphism θ . Let $u = t^{\frac{e}{m}}$, and consider the following $\mathbb{Z}$ -graded Lie algebra

$$\mathfrak{g}(\theta, m) := \bigoplus_{i \in \mathbb{Z}} u^i \mathfrak{g}_i \subset \mathfrak{g}[u, u^{-1}],$$

with $\mathfrak{g}(\theta, m)_i := u^i \mathfrak{g}_i$. In his book, Kac proved the following.

Theorem 3.3 (See [Kac], Theorem 8.5).

- (1) Let $\check{\lambda} = mx \in \check{X}^{\sigma}$. The automorphism $\text{Ad}(\lambda(u^{-1})) : \mathfrak{g}[u, u^{-1}] \simeq \mathfrak{g}[u, u^{-1}]$ induces an isomorphism

$$\Phi : \mathfrak{g}(\theta, m) \simeq {}^{\sigma}\tilde{\mathfrak{g}}.$$

- (2) Under the isomorphism Φ , the $\mathbb{Z}$ -grading of $\mathfrak{g}(\theta, m)$ becomes to the Kac-Moy-Prasad grading of ${}^{\sigma}\tilde{\mathfrak{g}}$ associated to x .
- (3) Under the isomorphism Φ , the derivation $u\partial_u$ on $\mathfrak{g}(\theta, m)$ becomes the derivation $D = \frac{m}{e}t\partial_t + \text{ad}\check{\lambda}$ on ${}^{\sigma}\tilde{\mathfrak{g}}$.
- (4) The invariant form $\langle u^i x, u^j y \rangle = \delta_{i,-j}(x, y)_{\text{Kill}}$ on $\mathfrak{g}[u, u^{-1}]$ induced an invariant from $\langle, \rangle_{\theta}$ (resp. $\langle, \rangle_{\sigma}$) on $\mathfrak{g}(\theta, m)$ (resp. ${}^{\sigma}\tilde{\mathfrak{g}}$) which is compatible with the grading, i.e., we have $\langle v, w \rangle_{\theta} = 0$ (resp. $\langle v, w \rangle_{\sigma} = 0$) for $v \in \mathfrak{g}(\theta, m)_i, w \in \mathfrak{g}(\theta, m)_j, i + j \neq 0$ (resp $v \in {}^{\sigma}\tilde{\mathfrak{g}}_{x,i}, w \in {}^{\sigma}\tilde{\mathfrak{g}}_{x,j}, i + j \neq 0$).

We have the following Corollary:

Corollary 3.4 ([Kac], [RY]). 1) For each $i = 0, 1, \dots, m-1$, there is a canonical isomorphism

$${}^{\sigma}\tilde{\mathfrak{g}}_{x,i} = \bigoplus_k t^{\frac{e(i-k)}{m}} \mathfrak{g}_i(k) \simeq \mathfrak{g}_i.$$

where the sum is over $-m + es_0 \leq k \leq m - es_0, k \equiv i \pmod{\frac{m}{e}}$.

2) If $i > 0$, all powers $t^{\frac{e(i-k)}{m}}$ appearing in the above sum are positive, i.e., for $0 < i < m, \mathfrak{g}_i(k) = 0$ unless $-m + es_0 \leq k \leq i, k \equiv i \pmod{\frac{m}{e}}$.

Proof. Since $\mathfrak{g}(\theta, m)_i = u^i \mathfrak{g}_i$ for $i = 0, 1, \dots, m-1$, result in §2.2 and above Theorem implies

$${}^\sigma \tilde{\mathfrak{g}}_{x,i} = \Phi(u^i \mathfrak{g}_i) = \text{Ad}(\lambda(u^{-1}))(u^i \mathfrak{g}_i) = \bigoplus_k t^{\frac{e(i-k)}{m}} \mathfrak{g}_i(k),$$

here the sum is over $-m + es_0 \leq k \leq m - es_0$, $k \equiv i \pmod{\frac{m}{e}}$. Part 1) follows. Since x is in the fundamental alcove $\bar{C}^\sigma$, direct calculation shows that, for $i > 0$, ${}^\sigma \tilde{\mathfrak{g}}_{x,i}$ is contained in ${}^\sigma \tilde{\mathfrak{g}} \cap \mathfrak{g}[t]$. Part 2) follows. □

Let us assume θ is regular and let $X \in \mathfrak{g}_1^r$ be a regular semi-simple element. Consider

$$p_1 = \Phi(uX) \in {}^\sigma \tilde{\mathfrak{g}}_{x,1}.$$

We have the following generalization of [Kac1, Proposition 3.8]

Proposition 3.5. *Let $\mathfrak{a} = \text{Ker}(\text{ad}p_1)$ and $\mathfrak{c} = \text{Im}(\text{ad}p_1)$. We have*

- (1) *The twisted loop algebra ${}^\sigma \tilde{\mathfrak{g}}$ has an orthogonal decomposition ${}^\sigma \tilde{\mathfrak{g}} = \mathfrak{a} \oplus \mathfrak{c}$ with respect to the invariant form $\langle, \rangle_\sigma$ in Theorem 3.3.*
- (2) *The Lie subalgebra $\mathfrak{a}$ is commutative. With respect to the Kac-Moy-Prasad grading, $\mathfrak{a} = \bigoplus \mathfrak{a}_i$, the subspaces $\mathfrak{a}_i$ and $\mathfrak{a}_j$ are orthogonal (resp. non degenerately paired) with respect to the invariant form $\langle, \rangle_\sigma$ on ${}^\sigma \tilde{\mathfrak{g}}$ if $i + j \neq 0$ (resp. $i + j = 0$).*
- (3) *Consider the Kac-Moody central extension ${}^\sigma \tilde{\mathfrak{g}} \oplus \mathbb{C}K$ of ${}^\sigma \tilde{\mathfrak{g}}$ (cf. §2.2). The pre-image $\hat{\mathfrak{a}} = \mathfrak{a} \oplus \mathbb{C}K$ of $\mathfrak{a}$ in ${}^\sigma \tilde{\mathfrak{g}} \oplus \mathbb{C}K$ is a non-split central extension of $\mathfrak{a}$.*

Proof. We first prove part (1) and (2). Since the isomorphism $\Phi : \mathfrak{g}(\theta, m) \simeq {}^\sigma \tilde{\mathfrak{g}}$ is compatible with the invariant forms on both side, it is enough to prove the corresponding statement for $\mathfrak{g}(\theta, m)$. Let $\mathfrak{s} = \bigoplus_{i \in \mathbb{Z}/m} \mathfrak{s}_i$ be the centralizer of X in $\mathfrak{g}$ and $\mathfrak{b} = \bigoplus_{i \in \mathbb{Z}/m} \mathfrak{b}_i$ be its orthogonal complement with respect to the Killing form $(,)_{Kill}$. Consider $\mathfrak{a}' = \text{Ker}(\text{ad}(uX))$ and its orthogonal complement $\mathfrak{c}'$ in $\mathfrak{g}(\theta, m)$ with respect to the invariant form $\langle, \rangle_\theta$. We have

$$\mathfrak{a}' = \bigoplus_{i \in \mathbb{Z}} \mathfrak{a}'_i, \quad \mathfrak{c}' = \bigoplus_{i \in \mathbb{Z}} \mathfrak{c}'_i,$$

where $\mathfrak{a}'_i = u^i \mathfrak{s}_i$, $\mathfrak{c}'_i = u^i \mathfrak{b}_i$. Now since the restriction of $(,)_{Kill}$ to any Cartan subalgebra is non-degenerate and $\text{ad}(X)$ is invertible on $\mathfrak{b}$, we have i) $\mathfrak{b}_{i+1} = [X, \mathfrak{b}_i]$ and ii) $\mathfrak{s}_i$ and $\mathfrak{s}_j$ are orthogonal (resp. non degenerately paired) with respect to $(,)_{Kill}$ if $i + j \neq 0$ (resp. $i + j = 0$). Part (1) and (2) follow.

For part (3), we first notice that a cocycle corresponding to the central extension is given by ${}^\sigma \tilde{\mathfrak{g}} \times {}^\sigma \tilde{\mathfrak{g}} \rightarrow \mathbb{C}$, $(v, w) \rightarrow \langle t\partial_t(v), w \rangle_\sigma$. Thus it is enough to show that for any $z \neq 0 \in \mathfrak{a}_n$ there exists $z' \in \mathfrak{a}_{-n}$ such that $\langle t\partial_t(z), z' \rangle_\sigma \neq 0$. Now observe that

$$\langle t\partial_t(z), z' \rangle_\sigma = \langle \frac{e}{m}(D - \text{ad}\check{\lambda})(z), z' \rangle_\sigma = \langle \frac{ne}{m}z, z' \rangle_\sigma - \langle \frac{e}{m}[\check{\lambda}, z], z' \rangle_\sigma,$$

here D is the derivation of ${}^\sigma\tilde{\mathfrak{g}}$ in Theorem 3.3. Since $\mathfrak{a}$ is commutative by part (2) we have $\langle \frac{e}{m}[\tilde{\lambda}, z], z' \rangle_\sigma = \langle \frac{e}{m}\tilde{\lambda}, [z, z'] \rangle_\sigma = 0$, hence $\langle t\partial_t(z), z' \rangle_\sigma = \langle \frac{ne}{m}z, z' \rangle_\sigma$ and the desired claim follows again from part (2). $\square$

4. YUN'S θ -CONNECTIONS

In [Yun1], Z. Yun associated to each torsion automorphism $\theta \in \text{Aut}(\mathfrak{g})$ and a nonzero vector $X \in \mathfrak{g}_1$, a *twisted* flat G -connection ∇^X on the trivial G -bundle on $\mathbb{G}_m = \mathbb{P}^1 - \{0, \infty\}$, called the θ -connection associated to X . In this section recall his construction of ∇^X .

4.1. Twisted flat G -connection and cohomological rigidity. In this subsection we recall the definition of twisted flat G -connection and cohomological rigidity (see [Yun2] in the setting of ℓ -adic sheaves). Let C be a smooth curve and $\tilde{C} \rightarrow C$ be a finite etale Galois cover with Galois group Γ . Let $\sigma : \Gamma \rightarrow \text{Aut}(G)$ be a homomorphism. We define a σ -*twisted* flat G -connection on C to be a triple $(\mathcal{F}, \nabla, \delta)$ where $\mathcal{F}$ is G -bundle on $\tilde{C}$, ∇ is a flat G -connection on it, and δ is a collection of isomorphisms $\delta_\gamma : (\mathcal{F}, \nabla) \simeq \gamma^*(\mathcal{F}, \nabla)$, $\gamma \in \Gamma$ satisfying the usual cocycle relations with respect to the multiplication on Γ .

When $\mathcal{F}$ is the trivial G -bundle, then a σ -twisted flat connection on it may be described as an operator

$$\nabla = d + A(t)dt,$$

where d is the exterior derivative and $A(t)dt$ is a Γ -invariant one form on $\tilde{C}$, with Γ acting by deck transformation and by the map $\Gamma \xrightarrow{\sigma} \text{Aut}(G) \rightarrow \text{Aut}(\mathfrak{g})$ on $\mathfrak{g}$.

Let $(\mathcal{F}, \nabla, \delta)$ be a σ -twisted flat G -connection on a smooth curve C . Then the corresponding flat vector bundle ∇^{Ad} on $\tilde{C}$ associated to the adjoint representation descends to C by Γ -equivariance. Let $\bar{\nabla}^{\text{Ad}}$ be the flat vector bundle on C after descent.

Definition 4.1. A σ -twisted flat G -connection $(\mathcal{F}, \nabla, \delta)$ over an open subset C of a complete smooth curve $\bar{C}$ is called *cohomologically rigid* if

$$H^*(\bar{C}, j_{!*}\bar{\nabla}^{\text{Ad}}) = 0,$$

where $j : C \hookrightarrow \bar{C}$ is the inclusion, and $j_{!*}\bar{\nabla}^{\text{Ad}}$ is the *non-derived* push forward of the D -module $\bar{\nabla}^{\text{Ad}}$ along j . We use the notation $j_{!*}$ because in this case $j_{!*}\bar{\nabla}^{\text{Ad}}$ is also the intermediate extension of the D -module $\bar{\nabla}^{\text{Ad}}$ to $\bar{C}$.

4.2. Construction of ∇^X . We preserve the setup in §2.2. Let $\theta = \exp(x) \rtimes \sigma \in \text{Aut}(\mathfrak{g}) = G \rtimes \text{Aut}(R, \Delta)$ be a torsion automorphism of $\mathfrak{g}$. Let $\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}/m} \mathfrak{g}_i$ be the corresponding grading. Let $X \in \mathfrak{g}_1$ and let us write $X = \sum X_k$, $X_k \in \mathfrak{g}_1(k)$ according

to (5). By Corollary 3.4, we have $X_k = 0$ unless $-m + es_0 \leq k \leq 1$ and $k \equiv 1 \pmod{\frac{m}{e}}$. Define

$$p_1 = \Phi(uX) = \sum X_k t^{\frac{e(1-k)}{m}} \in {}^\sigma \tilde{\mathfrak{g}},$$

here Φ is the isomorphism in Theorem 3.3. Then the θ -connection associated to X is the following flat G -connection on the trivial G -bundle on $\mathbb{G}_m = \text{Spec} \mathbb{C}[t, t^{-1}]$

$$(7) \quad \nabla^X = d + p_1 \frac{dt}{t} = d + \sum_{-m+es_0 \leq k \leq 1, k \equiv 1 \pmod{\frac{m}{e}}} X_k t^{\frac{e(1-k)}{m}} \frac{dt}{t}.$$

Note that $\frac{e(i-k)}{m} \in \mathbb{Z}$.

The $\mathfrak{g}$ -valued one form $p_1 \frac{dt}{t}$ is σ -invariant, where σ acts on $\mathbb{G}_m$ by the formula $t \rightarrow \xi_e^{-1} t$ and by the pinned automorphism on $\mathfrak{g}$. Therefore, by the discussion in §4.1, we can regard ∇^X as a σ -twisted flat G -connection on the trivial G -bundle on $\mathbb{G}_m$, where we regard σ as a map $\sigma : \mu_e = \langle \xi_e \rangle \rightarrow \text{Aut}(G)$ sending $\xi_e \rightarrow \sigma$, and $\Gamma = \mu_e$ is the Galois group of the finite étale Galois cover $[e] : \tilde{\mathbb{G}}_m = \mathbb{G}_m \rightarrow \mathbb{G}_m$ given by the e -th power map. We will give several examples of θ -connections in §7.

4.3. Residue at 0. Notice that $\frac{e(1-k)}{m} > 0$ for $k < 1$, thus the connection ∇^X has regular singularity at 0 with residue $\text{Res}(\nabla^X) = X_1 \in \mathfrak{g}_1(1)^\sigma$. Moreover, since $\mathfrak{g}(1)^\sigma$ consists of nilpotent elements of $\mathfrak{g}^\sigma$, the residue is nilpotent.

There is a dense open subset of $\mathfrak{g}(1)^\sigma$ which lies in a single nilpotent G^σ -orbit of $\mathfrak{g}^\sigma$. We denote this orbit by $\mathcal{O}_\theta$. Thus, for generic $X \in \mathfrak{g}_1$ the residue $\text{Res}(\nabla^X)$ lies in $\mathcal{O}_\theta$.

The assignment $\theta \rightarrow \mathcal{O}_\theta$ gives a well defined map ²

$$\{\text{torsion automorphism of } \mathfrak{g} \text{ whose image in } \text{Aut}(R, \Delta) \text{ is } \sigma\} \rightarrow \{\text{nilpotent orbits in } \mathfrak{g}^\sigma\}.$$

We now assume θ is stable. Consider the normalized Kac coordinates $\{s_0, s_1, \dots, s_{\ell_\sigma}\}$ of θ . If we omit s_0 and double the remaining Kac coordinates we obtain the weighted Dynkin diagram for the nilpotent orbit $\mathcal{O}_\theta$. Thus for Y in $\mathcal{O}_\theta$, we have $\dim \mathfrak{g}^{\sigma, Y} = \dim \mathfrak{g}(0)^\sigma$. The nilpotent class $\mathcal{O}_\theta$ is distinguished if and only if $\dim \mathfrak{g}(0)^\sigma = \dim \mathfrak{g}(1)^\sigma$.

Recall that stable torsion automorphisms θ are classified by regular elliptic W -conjugacy classes in the coset $W\sigma$ (cf. [RLYG, Corollary 15]). We therefore get a map

$$(8) \quad \{\text{regular elliptic classes in } W\sigma\} \rightarrow \{\text{nilpotent orbits in } \mathfrak{g}^\sigma\}.$$

In the case $\sigma = \text{id}$ and the normalized Kac coordinates satisfies $s_0 = 1$, this map is studied in [S] and [RLYG, §7.3] (see §7.1 for more details). The relation between this map and Kazhan-Lusztig map [KL] is discussed in [RLYG, §8.3, Remark 2].

²This map and the map in (8) are due to Z. Yun.

We expect that for any stable vector $X \in \mathfrak{g}_1^s$ we have $X_1 \in \mathcal{O}_\theta$. In other words, we expect the conjugacy classes of the residue $\text{Res}(\nabla^X)$, $X \in \mathfrak{g}_1^s$ depends only on θ and is given by the map (8). We will verify this expectation in some examples in §7.

Remark 4.2. Recall that Heisenberg algebras of the Kac-Moody central extension ${}^\sigma \tilde{\mathfrak{g}} \oplus \mathbb{C}K$ are parametrized, up to conjugacy, by W -conjugacy classes of the coset $W\sigma$ (see, e.g., [KP] for the case $\sigma = \text{id}$). Given $w \in W\sigma$, let $\hat{\mathfrak{a}}_w$ denote the associated Heisenberg subalgebra. One can show that, when θ is stable torsion automorphism, the algebra $\hat{\mathfrak{a}}$ in Proposition 3.5 is conjugate to the Heisenberg sub-algebra $\hat{\mathfrak{a}}_w$ where w is an element in the regular elliptic conjugacy class of $W\sigma$ corresponding to θ .

4.4. Slopes and Irregularity at ∞ . Let us compute the slopes of ∇^X at ∞ . Consider the covering given by $t = a^{-\frac{m}{e}}$. Then the connection ∇^X becomes

$$d - \frac{m}{e} \sum_k X_k a^{k-1} \frac{da}{a}.$$

Taking the gauge transform with $\check{\lambda}^{-1}(a)$, then $\text{Ad}(\check{\lambda}^{-1}(a))X_i = a^{-k}X_k$, hence the connection becomes

$$(9) \quad d - \frac{m}{e} X \frac{da}{a^2} + \check{\lambda} \frac{da}{a}.$$

When X is semi-simple, using the definition of slopes in [FG, §5] (or [CK, §2.2]), we see that the slopes of the connection at ∞ are either 0 or e/m .

Recall that any representation V of G gives rise to a flat vector bundle $\nabla^{X,V}$ on $\mathbb{G}_m$. We compute the irregularity of the connection $\text{Irr}_\infty(\nabla^{X,V})$ at infinity when X is semi-simple following [FG, §13]. Since X is semi-simple the leading term of the connection in (9) is diagonalizable in any representation V of G . It implies the slopes of the connection $\nabla^{X,V}$ is either 0 or e/m , the former occurring at the zero eigenspaces of X on V and the later occurring at the non-zero eigenspaces. According to [Katz, §1 and §2.3], the irregularity $\text{Irr}_\infty(\nabla^{X,V})$ is equal to the sum of the slopes of the connection at ∞ . This implies

$$\text{Irr}_\infty(\nabla^{X,V}) = \frac{e}{m}(\dim V - \dim V^X).$$

Assume $\nabla^{X,V}$ descends to a flat vector bundle $\bar{\nabla}^{X,V}$ via the e -th power map $[e] : \tilde{\mathbb{G}}_m \rightarrow \mathbb{G}_m$. Then again by [Katz, §2.3], we have

$$(10) \quad \text{Irr}_\infty(\bar{\nabla}^{X,V}) = \text{Irr}_\infty(\nabla^{X,V})/e = \frac{1}{m}(\dim V - \dim V^X).$$

5. MAIN RESULT

We preserve the setup of §4.1. Let $\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}/m\mathbb{Z}} \mathfrak{g}_i$ be a grading of $\mathfrak{g}$ and let $\theta = \theta' \rtimes \sigma \in \text{Aut}(\mathfrak{g})$ be the corresponding automorphism. Let $X \in \mathfrak{g}_1$ be a nonzero vector

and ∇^X be the corresponding θ -connection, which is a σ -twisted flat G -connection on $\mathbb{G}_m$.

Consider the adjoint representation Ad of G on its Lie algebra $\mathfrak{g}$. The corresponding flat vector bundle $\nabla^{X, \text{Ad}}$ descends via the e -th power map $[e] : \widetilde{\mathbb{G}}_m \rightarrow \mathbb{G}_m$ by σ -equivariance. Let $\bar{\nabla}^{X, \text{Ad}}$ be the connection after descent.

Here are the main results of this note, which generalize [FG, Theorem 1 and Proposition 11] to general θ -groups:

Theorem 5.1. *Assume θ is regular. Then for any regular semi-simple vector $X \in \mathfrak{g}_1^r$, we have*

$$H^0(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = H^2(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = 0$$

and

$$(11) \quad \dim H^1(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = \frac{\#R}{m} - \dim \mathfrak{g}^{\sigma, X_1}.$$

Here $j : \mathbb{G}_m \hookrightarrow \mathbb{P}^1$ is the canonical embedding and $X_1 \in \mathfrak{g}_1(1)^\sigma$ is the residue of the connection ∇^X at 0 (see §4.3).

Theorem 5.2. *Assume θ is stable and its normalized Kac coordinates satisfies $s_0 = 1$. Then for any stable vector $X \in \mathfrak{g}_1^s$, we have*

$$H^i(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = 0$$

for all i . I.e., ∇^X is cohomologically rigid (see Definition 4.1).

6. PROOFS

6.1. The key step leading the proofs of Theorem 5.1 and Theorem 5.2 is the computation of the cohomology groups $H^0(D_0^\times, \bar{\nabla}^{X, \text{Ad}})$ and $H^0(D_\infty^\times, \bar{\nabla}^{X, \text{Ad}})$. Here $D_0^\times = \text{Spec} \mathbb{C}((t))$ (resp. $D_\infty^\times = \text{Spec} \mathbb{C}((t^{-1}))$) is the formal punctured disc around 0 (resp. ∞).

We first introduce some auxiliary notations that will be used in the rest of the section. Let $\nabla^X = d + p_1 \frac{dt}{t}$ be the θ -connection associated to $X \in \mathfrak{g}_1^s$ (see §4.2). Recall $p_1 = \sum X_k t^{\frac{e(1-k)}{m}} \in {}^\sigma \tilde{\mathfrak{g}}$. The connection ∇^X gives a $\mathbb{C}$ -linear map

$$\nabla^{X, \text{Ad}} : \mathfrak{g}[[t, t^{-1}]] \rightarrow \mathfrak{g}[[t, t^{-1}]] \frac{dt}{t}.$$

Let $f = \sum v_n t^n \in \mathfrak{g}[[t, t^{-1}]]$ be a solution to $\nabla^{X, \text{Ad}}(f) = 0$. The components v_n satisfy

$$(12) \quad nv_n + [X_1, v_n] + \sum_{-m+es_0 \leq i \leq 0, i \equiv 1 \pmod{\frac{m}{e}}} [X_i, v_{a_i}] = 0,$$

where $a_i = n - \frac{e(1-i)}{m} \in \mathbb{Z}$. Notice that $a_i < n$ for all i .

6.1.1. We compute $H^0(D_0^\times, \bar{\nabla}^{X, \text{Ad}})$. Recall $\bar{\nabla}^{X, \text{Ad}}$ is the descent of $\nabla^{X, \text{Ad}}$ along the e -th power map $[e] : \mathbb{G}_m \rightarrow \mathbb{G}_m$. Thus $H^0(D_0^\times, \bar{\nabla}^{X, \text{Ad}}) = H^0(D_0^\times, [e]^* \bar{\nabla}^{X, \text{Ad}})^\sigma = H^0(D_0^\times, \nabla^{X, \text{Ad}})^\sigma = \text{Ker}(\nabla^{X, \text{Ad}} : \mathfrak{g}((t))^\sigma \rightarrow \mathfrak{g}((t))^\sigma \frac{dt}{t})$. Let $f = \sum v_n t^n \in \mathfrak{g}((t))^\sigma$ be a solution to $\nabla^{X, \text{Ad}}(f) = 0$. If $f \neq 0$, then there exists $b \in \mathbb{Z}$ such that $v_b \neq 0$ and $v_s = 0$ for $s < b$. We claim that $b \geq 0$. Indeed, if $b < 0$, then equation (12) implies

$$bv_b + [X_1, v_b] = 0,$$

which is impossible since the operator $b \cdot \text{Id} + X_1$ is invertible (recall $X_1 \in \mathfrak{g}(1)^\sigma$ is nilpotent). Thus $f = \sum v_n t^n \in \mathfrak{g}[[t]]^\sigma$ and $v_0 \in \mathfrak{g}^\sigma$ lies in the kernel of $\text{ad}(X_1)$. The equation (12) also implies there is a unique solution $f = \sum v_n t^n$ in $\mathfrak{g}[[t]]^\sigma$ for each $v_0 \in \mathfrak{g}^{\sigma, X_1}$. Above discussion shows that

$$H^0(D_0^\times, \bar{\nabla}^{X, \text{Ad}}) = \text{Ker}(\nabla^{X, \text{Ad}} : \mathfrak{g}((t))^\sigma \rightarrow \mathfrak{g}((t))^\sigma \frac{dt}{t}) = \mathfrak{g}^{\sigma, X_1}.$$

6.1.2. We show that $H^0(D_\infty^\times, \bar{\nabla}^{X, \text{Ad}})$ is zero. For this, we need some preliminary results about solutions $f \in \mathfrak{g}[[t, t]]^\sigma$ to $\nabla^{X, \text{Ad}}(f) = 0$. Let f be such a solution. If we write f in its components for the *Kac-Moy-Prasad* grading: $f = \sum y_n$ where $y_n \in {}^\sigma \tilde{\mathfrak{g}}_{x, n}$, then we have

$$(\frac{m}{e} t \nabla^{X, \text{Ad}})(f) = \sum_n ((\frac{m}{e} t \partial_t + \text{ad} \check{\lambda}) y_n + \frac{m}{e} [p_1, y_n] - [\check{\lambda}, y_n]) dt = 0.$$

Recall $p_1 = \sum X_k t^{\frac{e(1-k)}{m}} \in {}^\sigma \tilde{\mathfrak{g}}$.

Notice that the operator $\frac{m}{e} t \partial_t + \text{ad} \check{\lambda}$ is exactly the derivation D of ${}^\sigma \tilde{\mathfrak{g}}$ in Theorem 3.3 which defines the *Kac-Moy-Prasad* grading. Thus we have $(\frac{m}{e} t \partial_t + \text{ad} \check{\lambda}) y_n = n y_n$ and above equation gives rise to the identity

$$(13) \quad n y_n - [\check{\lambda}, y_n] + \frac{m}{e} [p_1, y_{n-1}] = 0$$

for all $n \in \mathbb{Z}$.

We have the following lemma

Lemma 6.1 ([FG], Lemma 6). *Suppose that y_n satisfying (13) and $y_n \in \mathfrak{a}_n$ for some n . Then $y_m = 0$ for all $m \leq n$.*

Proof. Assume that $y_n \neq 0$. In the course of the proof of Corollary 3.5 (part (3)), we have shown that there exists $z \in \mathfrak{a}_{-n}$ such that $\langle t \partial_t(y_n), z \rangle_\sigma \neq 0$. On the other hand, since y_n satisfies (13), we have

$$t \partial_t(y_n) = \frac{e}{m} (D - \text{ad} \check{\lambda})(y_n) = \frac{e}{m} (n y_n - [\check{\lambda}, y_n]) = -[p_1, y_{n-1}] \in \mathfrak{c}$$

and it implies $\langle t \partial_t(y_n), z \rangle_\sigma = 0$. We get a contradiction. Hence y_n must be zero.

Now, the equation (13) shows that if $y_n = 0$ then $y_{n-1} \in \mathfrak{a}_{n-1}$, hence, by induction that $y_m = 0$ for all $m \leq n$. $\square$

Above lemma implies $H^0(D_\infty^\times, \bar{\nabla}^{X, \text{Ad}}) = 0$. To see this, observe that

$$H^0(D_\infty^\times, \bar{\nabla}^{X, \text{Ad}}) = \text{Ker}(\nabla^{X, \text{Ad}} : \mathfrak{g}((t^{-1}))^\sigma \rightarrow \mathfrak{g}((t^{-1}))^\sigma \frac{dt}{t}).$$

Let $f \in \text{Ker}(\nabla^{X, \text{Ad}} : \mathfrak{g}((t^{-1}))^\sigma \rightarrow \mathfrak{g}((t^{-1}))^\sigma \frac{dt}{t})$. Then we have $v_n = 0$ for $n \gg 0$. This implies $y_n = 0$ for $n \gg 0$ (recall that y_n are the components of f for the *Kac-Moy-Prasad* grading), hence by above lemma $y_n = 0$ for all n . So we must have $f = 0$.

6.2. Proof of Theorem 5.1. According to [FG, §8], we have

$$H^0(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = H^0(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}),$$

$$H^2(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = H_c^2(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}),$$

and there is an exact sequence

$$0 \rightarrow H^0(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) \rightarrow H^0(D_0^\times, \bar{\nabla}^{X, \text{Ad}}) \oplus H^0(D_\infty^\times, \bar{\nabla}^{X, \text{Ad}}) \rightarrow H_c^1(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) \rightarrow H^1(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) \rightarrow 0.$$

We first prove $H^0(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = H^2(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = 0$. Since $H^0(D_\infty^\times, \bar{\nabla}^{X, \text{Ad}}) = 0$ by the result in §6.1.2, $\bar{\nabla}^{X, \text{Ad}}$ admits no global sections, i.e., $H^0(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = H^0(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) = 0$. Dually, $H^2(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = H_c^2(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) = H^0(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}})^* = 0$. Here we used the fact adjoint representation Ad is self-dual, hence $(\bar{\nabla}^{X, \text{Ad}})^* \simeq \bar{\nabla}^{X, \text{Ad}}$.

Now we prove $\dim H^1(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = \#R/m - \dim \mathfrak{g}^{\sigma, X_1}$. Results from §6.1.1, §6.1.2 and above exact sequence imply

$$(14) \quad 0 \rightarrow \mathfrak{g}^{\sigma, X_1} \rightarrow H_c^1(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) \rightarrow H^1(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) \rightarrow 0.$$

Thus it suffices to prove that $\dim H_c^1(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) = \#R/m$.

Recall the Deligne's formula in [D, §6.21.1] for the Euler characteristic

$$\chi_c(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) := \sum_i (-1)^i \dim H_c^i(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) = \chi_c(\mathbb{G}_m) \text{rank}(\bar{\nabla}^{X, \text{Ad}}) - \sum_{\alpha=0, \infty} \text{Irr}_\alpha(\bar{\nabla}^{X, \text{Ad}}).$$

Since $\chi_c(\mathbb{G}_m) = 0$ and $\bar{\nabla}^{X, \text{Ad}}$ is regular at 0, it implies

$$\chi_c(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) = -\text{Irr}_\infty(\bar{\nabla}^{X, \text{Ad}}).$$

Using the vanishing of H_c^0 , H_c^2 and the formula in line (10), we get

$$\dim H_c^1(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) = \text{Irr}_\infty(\bar{\nabla}^{X, \text{Ad}}) = \frac{1}{m}(\dim \mathfrak{g} - \dim \mathfrak{g}^X).$$

Since X is regular semi-simple, we have $\frac{1}{m}(\dim \mathfrak{g} - \dim \mathfrak{g}^X) = \#R/m$, hence

$$\dim H_c^1(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) = \#R/m.$$

This finished the proof of Theorem 5.1.

6.3. Proof of Theorem 5.2. It is enough to show that $H^1(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}}) = 0$. We begin with the following lemma:

Lemma 6.2. *For any solution $f = \sum v_n t^n$ of $\nabla^{X, \text{Ad}}(f) = 0$ in $\mathfrak{g}[[t, t^{-1}]]^\sigma$ we have $v_n = 0$ for all $n < 0$.*

Proof. Write $f = \sum y_n$ in the components for the *Kac-Moy-Prasad* grading. When $n = 0$, the equation (13) becomes

$$-[\check{\lambda}, y_0] + m[p_1, y_{-1}] = 0.$$

Since $s_0 = 1$ by assumption, Lemma 2.2 implies $y_0 \in \mathfrak{g}_0 = \mathfrak{g}_0 \cap \mathfrak{g}(0) \subset \ker(\text{ad } \check{\lambda})$. Therefore above equation implies $y_{-1} \in \mathfrak{a}_{-1}$, thus by Lemma 6.1, we have $y_n = 0$ for $n < 0$, or equivalently $f = \sum_{n \geq 0} y_n$. On the other hand, Corollary 3.4 and the fact $\mathfrak{g}_0 = \mathfrak{g}_0 \cap \mathfrak{g}(0)$ imply $y_n \in \mathfrak{g}[t]$ for $n \geq 0$. The Lemma follows. $\square$

By [FG, §9], we have $H_c^1(\mathbb{G}_m, \bar{\nabla}^{X, \text{Ad}}) \simeq \text{Ker}(\nabla^{X, \text{Ad}} : \mathfrak{g}[[t, t^{-1}]]^\sigma \rightarrow \mathfrak{g}[[t, t^{-1}]]^\sigma \frac{dt}{t})$, which is equal to $\text{Ker}(\nabla^{X, \text{Ad}} : \mathfrak{g}[[t]]^\sigma \rightarrow \mathfrak{g}[[t]]^\sigma \frac{dt}{t})$ by above Lemma. The same argument as in §6.1.1 shows that $\text{Ker}(\nabla^{X, \text{Ad}} : \mathfrak{g}[[t]]^\sigma \rightarrow \mathfrak{g}[[t]]^\sigma \frac{dt}{t}) = \mathfrak{g}^{\sigma, X_1}$. Therefore the first two terms in the short exact sequence (14) both have dimension $\dim \mathfrak{g}^{\sigma, X_1}$. This proves the vanishing of $H^1(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \text{Ad}})$, hence finished the proof of Theorem 5.2

7. EXAMPLES

In this section we give several examples of θ -connections ∇^X . In each example we write down the connections explicitly and check its cohomological rigidity using the formula in (11). We also check that, in each case, the residue of the θ -connection at $0 \in \mathbb{P}^1$ (or rather its conjugacy classes) depends only on θ , hence verify our expectation in §4.3. References for this section are [FG, RLYG, RY].

7.1. S-distinguished nilpotent case. Recall that a nilpotent element N in $\mathfrak{g}$ is called *distinguished* if $\mathfrak{g}^N$ consists of nilpotent elements. Let $N \in \mathfrak{g}$ be a distinguished nilpotent element. There is a co-character $\check{\lambda}$ such that $\text{Ad } \check{\lambda}(t)N = tN$ for all $t \in \mathbb{C}^\times$. This gives a grading $\mathfrak{g} = \bigoplus_{k=-a}^a \mathfrak{g}(k)$ where $\mathfrak{g}(k) = \{x \in \mathfrak{g} | \text{Ad } \check{\lambda}(t)x = t^k x\}$. Set $m = a + 1$ and consider the inner automorphism $\theta_N := \check{\lambda}(\xi_m) \in \text{Aut}(\mathfrak{g})$. We have $\mathfrak{g}_0 = \mathfrak{g}(0)$ and $\mathfrak{g}_1 = \mathfrak{g}(1) \oplus \mathfrak{g}(-a)$. Following [RLYG, §7.3], we say that a distinguished nilpotent element $N \in \mathfrak{g}$ is *S-distinguished* if the automorphism θ_N is stable. According to *loc. cit.*, a nilpotent element N is S-distinguished if and only if there exists $E \in \mathfrak{g}(-a)$ such that $N + E \in \mathfrak{g}_1$ is stable. Moreover, assume $\mathfrak{g}$ is of exceptional type, the map $N \rightarrow \theta_N$ defines a bijection between the set of S-distinguished nilpotent orbits in $\mathfrak{g}$ to the set of stable inner automorphism on $\mathfrak{g}$ with $s_0 = 1$.

Let θ_N be the stable automorphism of $\mathfrak{g}$ corresponding to a S-distinguished nilpotent element $N \in \mathfrak{g}$. Let $X = N + E \in \mathfrak{g}_1 = \mathfrak{g}(1) \oplus \mathfrak{g}(-a)$ be a stable vector. Then

by the formula in (7), the corresponding θ -connection ∇^X takes the form

$$\nabla^X = d + \frac{N}{t}dt + E dt.$$

Note that N is the residue of ∇^X at zero.

Let us verify that ∇^X is cohomologically rigid, i.e., $\dim H^*(\mathbb{P}^1, j_{!*}\nabla^{X, \text{Ad}}) = 0$. By Theorem 5.1, we have $H^0 = H^2 = 0$. Thus it remains to show $H^1(\mathbb{P}^1, j_{!*}\nabla^{X, \text{Ad}}) = \frac{\#R}{m} - \dim \mathfrak{g}^N = 0$. To see this recall that N is distinguished, thus we have $\dim \mathfrak{g}^N = \dim \mathfrak{g}(0) = \dim \mathfrak{g}_0$. On the other hand, we have $\dim \mathfrak{g}_0 = \frac{\#R}{m}$ (see [P, Theorem 4.2]). Result follows.

7.1.1. Frenkel-Gross case. We preserve the setup in §2.2. Consider the regular nilpotent element $N = \sum_{i=1}^l E_i$ in $\mathfrak{g}$. By [Kos, Corollary 6.4], it is S-distinguished. We can take the co-character to be $\check{\lambda} = \check{\rho}$ and $\theta_N = \check{\rho}(\xi_h) \in \text{Aut}(\mathfrak{g})$, where $\check{\rho}$ is the half-sum of positive co-roots and h is the Coxeter number.

We have $\mathfrak{g}_0 = \mathfrak{t}$ and $\mathfrak{g}_1 = \mathfrak{g}(1) \oplus \mathfrak{g}(-h+1)$, $\mathfrak{g}(1) = \bigoplus_{i=1}^\ell \mathfrak{g}_{\alpha_i}$, $\mathfrak{g}(-h+1) = \mathfrak{g}_{-\beta}$. Here β is the highest root. Choosing a generator E_0 for $\mathfrak{g}_{-\beta}$ and identifying $\mathfrak{g}_1$ with $\bigoplus_{i=0}^\ell \mathbb{C}E_i$, the open subset $\mathfrak{g}_1^s$ of stable vectors can be identified with $\mathfrak{g}_1^s = \{\sum c_i E_i | c_i \neq 0 \text{ for } i = 0, \dots, \ell\}$. For any $X = \sum c_i E_i \in \mathfrak{g}_1^s$, the corresponding θ -connection takes the form

$$\nabla^X = d + \frac{\sum_{i=1}^\ell c_i E_i}{t} dt + c_0 E_0 dt.$$

This is the rigid connections constructed in [FG]. Note that the residue of ∇^X at 0 is $N' = \sum_{i=1}^\ell c_i X_i$, which is regular nilpotent.

7.1.2. Type G_2 . Let $\mathfrak{g}$ is the simple Lie algebra of type G_2 . Let α_1, α_2 be the simple root of $\mathfrak{g}$, where α_2 is the short root. Consider the automorphism $\theta = \check{\lambda}(\xi_3)$, where $\check{\lambda} = \check{\omega}_1$ is the fundamental co-weight dual to α_1 and ξ_3 is a 3-th primitive root of unity. According to [RLYG], θ is a stable inner automorphism of order 3 with normalized Kac coordinates

$$1 \ 1 \Rightarrow 0.$$

Observe that if we omit s_0 and double remaining the Kac coordinates we obtain

$$2 \Rightarrow 0,$$

which is the weighted Dynkin diagram for the nilpotent orbit $G_2(2)$. This implies θ is equal to θ_N in §7.1 for some $N \in G_2(2)$.

We have $G_0 = \text{GL}_2(\mathbb{C})$ and $\mathfrak{g}_1 = \mathfrak{g}(1) \oplus \mathfrak{g}(-2)$, $\mathfrak{g}(1) = \bigoplus_{k=0}^3 \mathfrak{g}_{\alpha_1 + k\alpha_2}$, $\mathfrak{g}(-2) = \mathfrak{g}_{-2\alpha_1 - 3\alpha_2}$. As a representation of $G_0 = \text{GL}_2(\mathbb{C})$, we have

$$(15) \quad \mathfrak{g}(1) \simeq \det^2 \otimes P_3, \quad \mathfrak{g}(-2) \simeq \det^{-1} \otimes P_0,$$

where P_d is the space of homogeneous polynomials of degree d on $\mathbb{C}^2$, with the natural action of $G_0 = \mathrm{GL}_2(\mathbb{C})$. Choosing coordinates, we regard a vector $X \in \mathfrak{g}_1$ as a pair (f, z) , where $f = f(x, y)$ is a binary cubic polynomials over $\mathbb{C}$ and $a \in \mathbb{C}$. According to [RY, §7.5], we have $(f, z) \in \mathfrak{g}_1^s$ if and only if $z \neq 0$ and f has three distinct roots in the projective line. For any $X \in \mathfrak{g}_1^s$, let us write $X = X_1 + X_{-2}$ according to the decomposition $\mathfrak{g}_1 = \mathfrak{g}(1) \oplus \mathfrak{g}(-2)$. The corresponding θ -connection takes the form

$$\nabla^X = d + \frac{X_1}{t} dt + X_{-2} dt.$$

We claim that the residue X_1 is in the subregular nilpotent orbit $G_2(2)$. In particular, the conjugacy classes of the residue $\mathrm{Res}(\nabla^X)$ is independent of the choice $X \in \mathfrak{g}_1^s$. To prove the claim, observe that the intersection of $G_2(2)$ with $\mathfrak{g}(1)$ is open dense. Thus to show that X_1 is in $G_2(2)$ it is enough to show that $\dim \mathrm{Ad}G_0(X_1) = \dim \mathfrak{g}(1) = 4$. But it follows from the fact that the centralizer $Z_{G_0}(X_1)$ of X_1 in G_0 is the symmetric group S_3 (permuting the roots of f , where f is the binary cubic polynomial corresponding to X_1 under the isomorphism (15)), hence $\dim \mathrm{Ad}G_0(X_1) = \dim G_0 = 4$.

7.2. Type ${}^2A_{2n}$. Let $\mathfrak{g} = \mathfrak{sl}_{2n+1}(\mathbb{C})$ ($n \geq 1$). Let σ be a pinned automorphism of $\mathfrak{g}$. We define $\theta = \check{\rho}(-1) \rtimes \sigma \in \mathrm{Aut}(\mathfrak{g})$. According to [RLYG], it is a stable involution with Kac coordinates

$$1 \Rightarrow 0 \ 0 \cdots 0 \ 0 \Rightarrow 0.$$

Let $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ be the corresponding grading. We give a description of $\mathfrak{g}_0$ and $\mathfrak{g}_1$. Let V be a vector space over $\mathbb{C}$ of dimension $2n+1$ with basis $\{x_{-n}, \dots, x_{-1}, x_0, x_1, \dots, x_n\}$. We define an inner product $\langle, \rangle$ on V by the formula $\langle \sum a_i x_i, \sum b_i x_i \rangle = \sum_{i=-n}^n a_i b_{-i}$. For any $X \in \mathfrak{gl}(V)$, let X^* be the adjoint of X with respect to this inner product. Then under the canonical isomorphism $\mathfrak{sl}(V) \simeq \mathfrak{g}$, we have $\theta(X) = -X^*$ for any $X \in \mathfrak{g}$. Thus $\mathfrak{g}_0 \simeq \mathfrak{so}(V) = \{X \in \mathfrak{sl}(V) | X = -X^*\}$, $\mathfrak{g}_1 = \{X \in \mathfrak{sl}(V) | X = X^*\}$. Moreover, we have $\mathfrak{g}_1^s = \mathfrak{g}_1 \cap \mathfrak{g}^{rs}$, here $\mathfrak{g}^{rs}$ is the open subset of regular semi-simple elements in $\mathfrak{g}$.

Since $m = e = 2$ (recall m and e are the order of θ and σ), Corollary 3.4 implies $\mathfrak{g}_1 = \mathfrak{g}_1(0)$. Thus for any $X \in \mathfrak{g}_1$, the θ -connection ∇^X has the form

$$\nabla^X = d + X dt.$$

In particular, it is unramified at zero.

Finally, since $\mathfrak{g}^\sigma$ is a simple lie algebra of type B_n we have $\dim \mathfrak{g}^\sigma = n(2n+1)$. Thus for $X \in \mathfrak{g}_1^s$ we have

$$\dim H^1(\mathbb{P}^1, j_{!*} \bar{\nabla}^{X, \mathrm{Ad}}) = \frac{\#R}{2} - \dim \mathfrak{g}^\sigma = \frac{2n(2n+1)}{2} - n(2n+1) = 0.$$

8. RELATION WITH [Yun] AND RAMIFIED GEOMETRIC LANGLANDS

Let G be a split simply-connected almost simple group over a finite field k . In [Yun], starting with a stable torsion automorphism $\theta = \theta' \rtimes \sigma \in \text{Aut}(\mathfrak{g})$ and a stable linear function $X \in \mathfrak{g}_1^{*,s}$, the author construct an σ -twisted ℓ -adic $\check{G}$ -local system on $\mathbb{G}_m$. One can carry out above construction over the complex number with ℓ -adic sheaves replaced by D -modules. Starting with a stable torsion automorphism θ of $\mathfrak{g}$ over $\mathbb{C}$ and a stable linear function $\phi \in \mathfrak{g}_1^{*,s}$, we get a σ -twisted flat $\check{G}$ -connection $\text{KL}_{\check{G}}(\phi)$ on $\mathbb{G}_m$. Fixing $\sigma \in \text{Aut}(R, \Delta)$, the stable torsion automorphism θ is completely determined by a number m , so that θ is G -conjugate to $\check{\rho}(\xi_m) \rtimes \sigma$. Consider the stable automorphism θ for the dual Lie algebra $\check{\mathfrak{g}}$ determined by σ and m .

Conjecture 8.1 ([Yun1] and [HNY] Conjecture 2.14). *There is a bijection between the set of stable linear functions $\phi \in \mathfrak{g}_1^{*,s}$ and the set of stable vectors $X \in \check{\mathfrak{g}}_1^s$, such that whenever ϕ corresponds to X under this bijection, there is a natural isomorphism between σ -twisted flat $\check{G}$ -connection on $\mathbb{G}_m$*

$$\text{KL}_{\check{G}}(\phi) \simeq \nabla^X.$$

When $m = h$ is the Coxeter number, i.e., in the Frenkel-Gross case (see §7.1.1), above conjecture was proved in [Zhu] using a ramified version of Beilinson-Drinfeld's work on quantization of Hitchin's integrable systems. With X. Zhu and M. Kamgarpour, we plan to extend the methods in [Zhu] to more general stable automorphisms.

REFERENCES

- [CK] T.H. Chen, M. Kamgarpour.: *Preservation of depths in local geometric Langlands*, arXiv:1404.0598.
- [D] P. Deligne.: *Equations différentielles a points singuliers réguliers*, Lecture Notes in Mathematics, Vol. 163, Springer-Verlag, Berlin, 1970.
- [FG] E. Frenkel, D. Gross .: *Rigid irregular connection on the projective line* , Ann. of Math. (2) 170 (2009), no. 3, 1469-1512.
- [HNY] J.Heinloth, B-C. Ngô, Z.Yun.: *Kloosterman sheaves for reductive groups*, Ann. of Math. (2) 177 (2013), no. 1, 241-310.
- [KL] D. Kazhdan, G. Lusztig .: *Fixed point varieties on affine flag manifolds*, Israel Journal of Mathematics 1988, Volume 62, Issue 2, 29-168
- [Kac] V. Kac .: *Infinite dimensional Lie algebras* , 3rd Edition, Cambridge University Press, 1990.
- [Kac1] V. Kac .: *Infinite-dimensional algebras, Dedekind's η -function, classical Möbius function and the very strange formula*, Adv. Math. 30 (1978) 85-136.
- [Kos] B. Kostant.: *The principal three-dimensional subgroups and the Betti numbers of a complex simple Lie group*, Amer J. Math. 81. 973-1032 (1959)
- [KP] V. Kac, D. Peterson.: *112 construction of the basic representation of the loop group of E_8* , Symposium on anomalies, geometry, topology (Chicago, Ill., 1985), 276-298, World Sci. Publishing, Singapore, 1985.
- [Katz] N. Katz.: *On the calculation of some differential Galois groups*, Invent. Math. 87 (1987), no. 1, 13-61.
- [OV] A.L. Onishchik. E.B. Vinberg.: *Lie Groups and Lie Algebra III*, Encyclopaedia of Mathematical Sciences, Vol. 41.

- [P] P. Panyushev.: *On invariant theory of θ -groups*, Jour. Algebra, 283 (2005), pp. 655-670.
- [RLYG] M. Reeder, P. Levy, J.K. Yu, B. Gross.: *Gradings of positive rank on simple Lie algebras*, Transformation Groups, 17, No. 4, (2012), 1123-1190.
- [RY] M. Reeder, J.-K. Yu.: *Epipelagic representations and invariant theory*, Journal of the AMS, Vol 27, Number 2, April 2014, 437-477.
- [S] T.A. Springer.: *Regular elements of finite reflection groups*, Invent. Math. 25 (1974), 159-198.
- [Yun] Z. Yun.: *Epipelagic representation and rigid local systems*, arXiv:1401.7647.
- [Yun1] Z. Yun.: *Note on θ -groups and connections*, unpublished note.
- [Yun2] Z. Yun.: *Rigidity in automorphic representations and local systems*, arXiv:1405.3035
- [Zhu] X. Zhu.: *Frenkel-Gross's irregular connection and Heinloth-Ngô-Yun's are the same*, arXiv:1210.2680

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, 2033 SHERIDAN ROAD, EVANSTON,
60208, USA

E-mail address: `chenth@math.northwestern.edu`